

First-order Predicate Logic
The Classical Decision Problem

Validity/satisfiability of arbitrary first-order formulas is undecidable.

What about subclasses of formulas?

Examples

$\forall x \exists y (P(x) \rightarrow P(y))$

Satisfiable? Resolution?

$\exists x \forall y (P(x) \rightarrow P(y))$

Satisfiable? Resolution?

The $\exists^*\forall^*$ class

Definition

The $\exists^*\forall^*$ class is the class of closed formulas of the form

$$\exists x_1 \dots \exists x_m \forall y_1 \dots \forall y_n F$$

where F is quantifier-free and contains no function symbols of arity > 0 .

This is also called the [Bernays-Schönfinkel class](#).

Corollary

Unsatisfiability is decidable for formulas in the \exists^\forall^* class.*

What if a formula is not in the $\exists^*\forall^*$ class?

Try to transform it into the $\exists^*\forall^*$ class!

Example

$$\forall y \exists x (P(x) \wedge Q(y))$$

Heuristic transformation procedure:

1. Put formula into NNF
2. Push all quantifiers into the formula as far as possible (“miniscoping”)
3. Pull out \exists first and \forall afterwards

Miniscoping

Perform the following transformations bottom-up,
as long as possible:

- ▶ $(\exists x F) \equiv F$ if x does not occur free in F
- ▶ $\exists x (F \vee G) \equiv (\exists x F) \vee (\exists x G)$
- ▶ $\exists x (F \wedge G) \equiv (\exists x F) \wedge G$ if x is not free in G
- ▶ $\exists x F$ where F is a conjunction,
 x occurs free in every conjunct,
and the DNF of F is of the form $F_1 \vee \dots \vee F_n$, $n \geq 2$:
 $\exists x F \equiv \exists x (F_1 \vee \dots \vee F_n)$

Together with the dual transformations for \forall

Example

$$\exists x (P(x) \wedge \exists y (Q(y) \vee R(x)))$$

Warning: Complexity!

The monadic class

Definition

A formula is **monadic** if it contains only unary (monadic) predicate symbols and no function symbol of arity > 0 .

Examples

All men are mortal. Sokrates is a man. Sokrates is mortal.

The monadic class is decidable

Theorem

Satisfiability of monadic formulas is decidable.

Proof Put into NNF. Perform miniscoping.

The result has no nested quantifiers (**Exercise!**).

First pull out all \exists , then all \forall .

Existentially quantify free variables.

The result is in the $\exists^*\forall^*$ class.

Corollary

Validity of monadic formulas is decidable.

The finite model property

Definition

A formula F has the **finite model property** (for satisfiability) if F has a model iff F has a finite model.

Theorem

If a formula has the finite model property, satisfiability is decidable.


Theorem

Monadic formulas have the finite model property.

Classification by quantifier prefix of prenex form

There is a **complete** classification of decidable and undecidable classes of formulas based on

- ▶ the form of the quantifier prefix of the prenex form
- ▶ the arity of the predicate and function symbols allowed
- ▶ whether “=” is allowed or not.



A complete classification

Only formulas without function symbols of arity > 0 ,
no restrictions on predicate symbols.

Satisfiability is decidable:

$\exists^* \forall^*$ (Bernays, Schönfinkel 1928)

$\exists^* \forall \exists^*$ (Ackermann 1928)

$\exists^* \forall^2 \exists^*$ (Gödel 1932)

Satisfiability is undecidable:

$\forall^3 \exists$ (Surányi 1959)

$\forall \exists \forall$ (Kahr, Moore, Wang 1962)

Why complete?

Famous mistake by Gödel: $\exists^* \forall^2 \exists^*$ with “=” is **undecidable**
(Goldfarb 1984)