

Propositional Logic Equivalences

Equivalence

Definition (Equivalence)

Two formulas F and G are (semantically) equivalent if $\mathcal{A}(F) = \mathcal{A}(G)$ for every assignment \mathcal{A} .

We write $F \equiv G$ to denote that F and G are equivalent.

Exercise

Which of the following equivalences hold?

$$(A \wedge (A \vee B)) \equiv A$$

$$(A \wedge (B \vee C)) \equiv ((A \wedge B) \vee C)$$

$$(A \rightarrow (B \rightarrow C)) \equiv ((A \rightarrow B) \rightarrow C)$$

$$(A \rightarrow (B \rightarrow C)) \equiv ((A \wedge B) \rightarrow C)$$

Observation

The following connections hold:

$$\begin{aligned} \models F \rightarrow G & \text{ iff } F \models G \\ \models F \leftrightarrow G & \text{ iff } F \equiv G \end{aligned}$$

NB: “iff” means “if and only if”

Reductions between problems (I)

- ▶ **Validity** to **Unsatisfiability** (and back):

$$\begin{aligned} F \text{ valid} & \text{ iff } \neg F \text{ unsatisfiable} \\ F \text{ unsatisfiable} & \text{ iff } \neg F \text{ valid} \end{aligned}$$

- ▶ **Validity** to **Consequence**:

$$F \text{ valid} \quad \text{iff} \quad \top \models F$$

- ▶ **Consequence** to **Validity**:

$$F \models G \quad \text{iff} \quad F \rightarrow G \text{ valid}$$

Reductions between problems (II)

- ▶ **Validity** to **Equivalence**:

$$F \text{ valid} \quad \text{iff} \quad F \equiv \top$$

- ▶ **Equivalence** to **Validity**:

$$F \equiv G \quad \text{iff} \quad F \leftrightarrow G \text{ valid}$$

Properties of semantic equivalence

- ▶ Semantic equivalence is an **equivalence relation** between formulas.
- ▶ Semantic equivalence is **closed under operators**:

If $F_1 \equiv F_2$ and $G_1 \equiv G_2$
then $(F_1 \wedge G_1) \equiv (F_2 \wedge G_2)$,
 $(F_1 \vee G_1) \equiv (F_2 \vee G_2)$ and
 $\neg F_1 \equiv \neg F_2$

Equivalence relation + Closure under Operations
=
Congruence relation

Replacement theorem

Theorem

Let $F \equiv G$. Let H be a formula with an occurrence of F as a subformula. Then $H \equiv H'$, where H' is the result of replacing an arbitrary occurrence of F in H by G .

Proof by induction on the structure of H .

Equivalences (I)

Theorem

$$\begin{aligned}(F \wedge F) &\equiv F \\ (F \vee F) &\equiv F\end{aligned}\quad (\text{Idempotence})$$

$$\begin{aligned}(F \wedge G) &\equiv (G \wedge F) \\ (F \vee G) &\equiv (G \vee F)\end{aligned}\quad (\text{Commutativity})$$

$$\begin{aligned}((F \wedge G) \wedge H) &\equiv (F \wedge (G \wedge H)) \\ ((F \vee G) \vee H) &\equiv (F \vee (G \vee H))\end{aligned}\quad (\text{Associativity})$$

$$\begin{aligned}(F \wedge (F \vee G)) &\equiv F \\ (F \vee (F \wedge G)) &\equiv F\end{aligned}\quad (\text{Absorption})$$

Equivalences (II)

$$(F \wedge (G \vee H)) \equiv ((F \wedge G) \vee (F \wedge H))$$

$$(F \vee (G \wedge H)) \equiv ((F \vee G) \wedge (F \vee H))$$

$$\neg\neg F \equiv F$$

$$\neg(F \wedge G) \equiv (\neg F \vee \neg G)$$

$$\neg(F \vee G) \equiv (\neg F \wedge \neg G)$$

$$\neg\top \equiv \perp$$

$$\neg\perp \equiv \top$$

$$(\top \vee G) \equiv \top$$

$$(\top \wedge G) \equiv G$$

$$(\perp \vee G) \equiv G$$

$$(\perp \wedge G) \equiv \perp$$

(Distributivity)

(Double negation)

(deMorgan's Laws)

Warning

The symbols \models and \equiv are **not** operators
in the language of propositional logic
but part of the meta-language for talking about logic.

Examples:

$\mathcal{A} \models F$ and $F \equiv G$ are not propositional formulas.
 $(\mathcal{A} \models F) \equiv G$ and $(F \equiv G) \leftrightarrow (G \equiv F)$ are nonsense.