First-Order Logic Compactness

[Harrison, Section 3.16]

More Herbrand Theory

Recall Gödel-Herbrand-Skolem:

Theorem

Let F be a closed formula in Skolem form. Then F is satisfiable iff its Herbrand expansion E(F) is (propositionally) satisfiable.

Can easily be generalized:

Theorem

Let S be a set of closed formulas in Skolem form. Then S is satisfiable iff E(S) is (propositionally) satisfiable.

Transforming sets of formulas

Recall the transformation of single formulas into equisatisfiable Skolem form: close, RPF, skolemize

Theorem

Let S be a countable set of closed formulas. Then we can transform it into an equisatisfiable set of closed formulas T in Skolem form.

We call this transformation function skolem.

- ► Can all formulas in *S* be transformed in parallel?
- Why "countable"?

Transforming sets of formulas

1. Put all formulas in S into RPF.

Problem in Skolemization step: How do we generate new function symbols if all of them have been used already in *S*?

2. Rename all function symbols in S: $f_i^k \mapsto f_{2i}^k$

The result: equisatisfiable countable set $\{F_0, F_1, \dots\}$.

Unused symbols: all f_{2i+1}^k

3. Skolemize the F_i one by one using the f_{2i+1}^k not used in the Skolemization of F_0, \ldots, F_{i-1}

Result is equisatisfiable with initial S.

Compactness

Theorem Let S be a countable set of closed formulas. If every finite subset of S is satisfiable, then S is satisfiable.